

**Название публикации:**

Oxidative functionalization of C-H compounds induced by the extremely efficient osmium catalysts (a review)

**Авторы:**

Shul'pin, GB [ 1,2 ] ; Vinogradov, MM [ 3 ] ; Shul'pina, LS [ 3 ]

[ 1 ] Russian Acad Sci, Semenov Inst Chem Phys, Dom 4, Moscow, Russia

[ 2 ] Plekhanov Russian Univ Econ, Dom 36, Moscow, Russia

[ 3 ] Russian Acad Sci, Nesmeyanov Inst Organoelement Cpds, Dom 28, Moscow, Russia

**Наименование журнала:**

CATALYSIS SCIENCE & TECHNOLOGY

Том: 8 Выпуск: 17 Стр.: 4287-4313

DOI: 10.1039/c8cy00659h

Опубликовано: SEP 7 2018

Тип документа: Review

**Аннотация:**

Osmium derivatives are not as popular among catalytic chemists as the compounds of iron (an analog of osmium), copper, or manganese. Although osmium is expensive, poisonous and volatile, it finds applications not only in cis-hydroxylation of olefins but its derivatives have recently been employed in the oxygenation of C-H compounds (saturated and aromatic hydrocarbons and alcohols) by hydrogen peroxide as well as organic peroxides. The turnover in alkane oxidations catalyzed by soluble osmium derivatives has been found to be much higher than analogous reactions that used complexes of other transition metals (e.g., iron, manganese or copper). Using certain additives, the authors of this review have developed osmium-containing systems that are extremely efficient in transforming alkanes into corresponding alkyl hydroperoxides, which can further be reduced to alcohols easily. The new catalytic systems described here are capable of producing ground-breaking results in the field of metal-complex catalysts.

**Ключевые слова:**

Hydrogen-peroxide oxidation; tert-butyl hydroperoxide; tolylimido rhenium(v) complexes; transition-metal-complexes; nonheme iron catalysts; fenton-like reactions; group-iii metals; selective oxidation; alkane oxidation; manganese complexes